


1 **What is claimed is**


2 1. Annonaceous acetogenins substantially pure compounds having the structures a-g.

3 a. muricin A having formula as:




6 wherein the muricin A having an α, β -unsaturated γ -lactone with a hydroxyl
7 group at C-4 position, a mono-THF ring placed between C-15 and C-18 with
8 one flanking hydroxyl in a threo conformation, two methylene groups of the
9 mono-THF ring corresponding to trans conformation, two hydroxyl groups at
10 C-26 and C-27 as vicinal diol assigned as threo based, and the stereochemistry
11 at C-34 on the γ -lactone fragment performed in (S)-configuration.

12 b. muricin B having formula as:



15 wherein the muricin B having an α, β -unsaturated γ -lactone with a hydroxyl
16 group at C-4 position, a mono-THF ring placed between C-15 and C-18 with
17 one flanking hydroxyl in a trans/threo conformation, two methylene groups of
18 the mono-THF ring corresponding to trans conformation, two hydroxyl groups
19 at C-26 and C-27 as vicinal diol assigned as threo based, and the
20 stereochemistry at C-34 on the γ -lactone fragment performed in (S)-
21 configuration.


22 c. muricin C having formula:



24 wherein the muricin C having an α, β -unsaturated γ -lactone with a hydroxyl


group at C-4 position, a mono-THF ring placed between C-17 and C-20 with one flanking hydroxyl in trans/threo or threo/trans conformation, two hydroxyl groups at C-24 and C-25 as vicinal diol assigned as threo based, and the stereochemistry at C-34 on the γ -lactone fragment performed in (S)-configuration.

d. muricin D having formula:




wherein the muricin D having an α, β -unsaturated γ -lactone with a hydroxyl group at C-4 position, a mono-THF ring placed between C-15 and C-18 with one flanking hydroxyl in threo/trans conformation, two hydroxyl groups at C-22 and C-23 as vicinal diol assigned as threo based.

e. muricin E having formula:



wherein the muricin E having an α, β -unsaturated γ -lactone with a hydroxyl group at C-4 position, a mono-THF ring placed between C-12 and C-15 with one flanking hydroxyl in threo/trans conformation, two hydroxyl groups at C-22 and C-23 as vicinal diol assigned as threo based.


f. muricin F having formula:



wherein the muricin F having an α, β -unsaturated γ -lactone with a hydroxyl group at C-4 position, a mono-THF ring placed between C-17 and C-20 with

one flanking hydroxyl in threo/trans conformation, two hydroxyl groups at C-27 and C-28 as vicinal diol assigned as threo based, and a double bond determined at C-24/C-25.

g. muricin G having formula:



wherein the muricin G having an α, β -unsaturated γ -lactone with a hydroxyl group at C-4 position, a mono-THF ring placed between C-16 and C-19 with one flanking hydroxyl in threo/trans/threo conformation, one hydroxyl groups formed at C-10, a double bond determined at C-23/C-24, and the stereochemistry at C-34 on the γ -lactone fragment performed in (S)-configuration.

2. A method for substantially purified extract of claim 1 from the species *Annona muricata*, wherein the method comprising:

extracting *Annona muricata* seeds repeatedly with MeOH at room temperature; evaporating and partitioning the combined MeOH extracts to yield CHCl_3 and aqueous extracts;

further separating the CHCl_3 layer into ten fractions by column chromatography on Si gel with gradient system of *n*-hexane- CHCl_3 and CHCl_3 -MeOH;

combining the eighth and ninth fractions together and then further separating into ten sub-fractions by column chromatography;

isolating and purifying the Annonaceous acetogenins compounds from the ten sub-fractions.

3. The method as claimed in claim 2 for substantially purified extract of claim 1

1 from the species *Annona muricata*, in which the muricin A (1), muricin B (2), muricin C
2 (3), and muricin F (6) are isolated and purified from the seventh sub-fraction by a
3 preparative reversed-phase method.

4 4. The method as claimed in claim 2 for substantially purified extract of claim 1
5 from the species *Annona muricata*, in which the muricin D (4), muricin E (5), and
6 muricin G (7) are isolated and purified from the eighth sub-fraction by a preparative
7 reversed-phase method.

8 5. An anti-tumor composition selectively comprising an amount of substantially
9 pure muricins of claim 1, wherein the muricins are effective and acted as an anti-tumor
10 agent and selectively combined with pharmaceutically acceptable salt, ester, and carrier
11 in the anti-tumor composition.

12 6. The annonaceous acetogenins compounds as claimed in claim 1, wherein the
13 substantially pure muricins are selectively used for the preparation of a pharmaceutical
14 composition for the treatment of a patient having a tumor.

15 7. The anti-tumor composition as claimed in claim 5, wherein the anti-tumor
16 composition is used for pharmaceutically treating a patient having hepatoma cancer.

17 8. A method of treating a patient having a tumor, wherein said method comprising
18 administering an effective amount of a pharmaceutical composition comprising
19 muricins of claim 1 to a patient afflicted with cancer.

20 9. A method for treating hepatoma cancer, said method comprising administering
21 to a patient afflicted with hepatoma cancer an effective amount of a pharmaceutical
22 composition comprising a substantially pure bioactive compound selected from the
23 group consisting of muricins of claim 1 and pharmaceutically acceptable salt, ester, or
24 carrier.